

Unified description of bulk and interface-enhanced spin pumping

S. M. Watts, J. Grollier,* C. H. van der Wal, and B. J. van Wees
*Physics of Nanodevices, Materials Science Center, University of Groningen,
 Nijenborgh 4, 9747 AG Groningen, The Netherlands*
 (Dated: February 2, 2008)

The dynamics of non-equilibrium spin accumulation generated in metals or semiconductors by rf magnetic field pumping is treated within a diffusive picture. The dc spin accumulation produced in a uniform system by a rotating applied magnetic field or by a precessing magnetization of a weak ferromagnet is in general given by a (small) fraction of $\hbar\omega$, where ω is the rotation or precession frequency. With the addition of a neighboring, field-free region and allowing for the diffusion of spins, the spin accumulation is dramatically enhanced at the interface, saturating at the universal value $\hbar\omega$ in the limit of long spin relaxation time. This effect can be maximized when the system dimensions are of the order of $\sqrt{2\pi D/\omega}$, where D is the diffusion constant. We compare our results to the interface spin pumping theory of A. Brataas *et al.* [Phys. Rev. B **66**, 060404(R) (2002)].

PACS numbers: 85.75.-d, 72.25.Mk, 75.70.Cn

A new direction in the field of spin electronics is the study of the interaction between spin currents and the magnetization dynamics of a ferromagnetic electrode into, or out of which the spin currents flow. In spin pumping devices [1, 2, 3], electron spins are pumped from a ferromagnet in resonance with an rf magnetic field, into an adjoining non-magnetic metal without accompanying charge currents. The so-called spin battery [2] is based on the interface spin-mixing conductance [4], which describes the coherent reflection and spin rotation within an exchange length (typically a few nm in conventional ferromagnets) of the interface. It has been experimentally verified by measuring the effect of adjoining metal layers on the Gilbert damping constant of a ferromagnetic layer [5, 6], due to the transfer of angular momentum away from the ferromagnetic and into adjacent metals [7]. For the spin battery, under specific conditions the universal value of $\hbar\omega$ is obtained for the spin accumulation, where ω is the precession frequency of the magnetization. On the other hand, a bulk-oriented approach based on mean-field theory [8] has been used to relate the intrinsic Gilbert damping to the spin relaxation processes resulting from the generation of non-equilibrium spin accumulation in the ferromagnet.

We present a model in which spin pumping with a rotating magnetic field can produce spin accumulation in bulk material. Moreover, in a hybrid system in which spins are allowed to diffuse from a "pumped" region into a neighboring region without fields, the spin accumulation can be dramatically enhanced near the interface. In both cases the spin accumulation reaches the universal limit $\hbar\omega$ for long spin relaxation times. Our approach is based on classical dynamics of spin ensembles in a diffusive system, and is applicable to spin pumping in metals and semiconductors driven by a rotating, applied magnetic field, as well as in weak ferromagnets driven by magnetization precession. However, we emphasize that our semi-classical approach cannot be applied directly to the con-

ventional strong ferromagnet regime in which the large exchange fields require a non-diffusive, quantum mechanical treatment [4] because of the very small magnetic coherence length, much less than the mean-free-path. The dc spin-accumulation induced by spin-pumping is first derived for a bulk, uniform system, for which we obtain the surprising result that bulk spin accumulation is generated even for small Zeeman splitting. Then, by connecting a magnetic field-free region and solving the boundary-value problem of diffusion across the interface, we find a transition from bulk to interface-enhanced spin accumulation.

We start with Bloch-type equations written in terms of the chemical potentials for the three spin directions, $\vec{f} = (f_x, f_y, f_z)$ that describe the space- and time-dependent, non-equilibrium spin accumulation $\vec{f}(x, t)$ (restricting the description to one spatial dimension x) [9]. The basis of our description is the following equation for the dynamics of $\vec{f}(x, t)$ in a rotating magnetic field $\vec{B}(x, t)$ [10, 11]:

$$-\frac{\partial \vec{f}}{\partial t} + \vec{I}(x, t) = -D\nabla^2 \vec{f} + \frac{\vec{f}}{\tau} + \frac{g\mu_B}{\hbar} \vec{B} \times \vec{f}, \quad (1)$$

where D is the diffusion constant and $\vec{I}(x, t)$ is a source term which will be described in the next paragraph. The length scales relevant to the problem are the spin diffusion length $\lambda = \sqrt{D\tau}$, the Larmor precession length $\lambda_B = \sqrt{2\pi D/|\vec{\omega}_B|}$, where $\hbar\vec{\omega}_B = g\mu_B\vec{B}$, and a length $\lambda_\omega = \sqrt{2\pi D/\omega}$ corresponding to the distance the spins can diffuse in one period $\frac{2\pi}{\omega}$. Eqn. 1, with the left hand side equal to zero, has been shown to give an accurate description of spin dynamics in metallic systems for time-independent magnetic fields [11, 12].

The key new ingredient of our approach is the source term:

$$\vec{I}(x, t) = -g\mu_B \frac{\partial}{\partial t} \vec{B}(x, t). \quad (2)$$




FIG. 1: (a) Magnetic field vector diagram. An applied rf field B_{xy} rotates with angular frequency ω in the x-y plane. A static magnetic field B_z is applied perpendicular to B_{xy} generating a field vector $\vec{B}$ that rotates around B_z with cone angle θ . (b) The general hybrid system with an interface at $x = 0$. The fields just described are applied in the "pumped" region I, $x < 0$. $\vec{B} = 0$ in region II, $x > 0$.

It describes the rate at which the oscillating magnetic field, via the Zeeman energy, pushes spins aligned with the magnetic field below the Fermi level, and anti-aligned spins above from the Fermi level. This mechanism can be seen as a source of locally-injected, time-dependent spin currents [13]. Without this source term, $\dot{\vec{f}} = 0$ in the steady state.

The role of Eqn. 2 is sketched out as follows: consider two magnetic fields applied in orthogonal directions, say B_x and B_y , equal in magnitude but oscillating out of phase by $\pi/2$: $B_x = B_{xy} \cos \omega t$ and $B_y = B_{xy} \sin \omega t$. The resulting I_x generates spin accumulation in f_x in phase with B_y , and subsequent precession yields spin accumulation in f_z . Likewise, I_y produces an f_y that is π out of phase with B_x , and subsequent precession yields spin accumulation in f_z with the same sign as for the I_x case. Both f_x and f_y oscillate about a mean of zero with frequency ω . However, f_z acquires a finite dc component. The steady-state solution of $\dot{\vec{f}}$ can be imagined as a spin accumulation vector that follows the rotating field vector, lagging by phase angles which depend on ω and τ .

Fig. 1a shows the basic field configuration we will consider: a rotating magnetic field B_{xy} and a perpendicular, static field B_z . The field vector is $\vec{B} = (B_{xy} \cos \omega t, B_{xy} \sin \omega t, B_z)$ in Cartesian coordinates. We first consider region I (Fig. 1b) alone; *i.e.*, a uniform (spatially invariant) system without boundaries. Dropping the diffusion term and rewriting Eqn.1 in a more concise form:

$$\tau \dot{\vec{f}} = -\vec{f} - \vec{\omega}_B \tau \times \vec{f} - \hbar \dot{\vec{\omega}}_B \tau, \quad (3)$$

where $\vec{\omega}_B = (\omega_{xy} \cos \omega t, \omega_{xy} \sin \omega t, \omega_z)$. Since the induced spin accumulations f_x and f_y will, in the steady state, oscillate with the same frequency as B_x and B_y , it will be advantageous to transform to a coordinate system in which the x and y axes rotate about the z axis with angular frequency $\vec{\omega} = \omega \hat{z}$, so that $\vec{B} = (B_{xy}, 0, B_z)$.

The time derivative of a vector $\vec{r}$ in the stationary frame transforms as $(\dot{\vec{r}})_{lab} \Rightarrow (\dot{\vec{r}})_{rot} + \vec{\omega} \times (\vec{r})_{rot}$. Using this relation, Eqn.3 now becomes

$$\tau \dot{\vec{f}} = -\vec{f} - (\vec{\omega}_B + \vec{\omega}) \tau \times \vec{f} - \hbar (\dot{\vec{\omega}}_B + \vec{\omega} \times \vec{\omega}_B) \tau, \quad (4)$$

In the last term, we have $\dot{\vec{\omega}}_B = 0$ in this frame and $\vec{\omega} \times \vec{\omega}_B = (0, \omega \omega_{xy}, 0)$. For the stationary analysis we set $\dot{\vec{f}} = 0$, and obtain the result [9]:

$$f_z = \frac{(\omega_{xy} \tau)^2}{1 + (\omega_{xy} \tau)^2 + ((\omega_z + \omega) \tau)^2} \hbar \omega. \quad (5)$$

The other components are related to f_z : $f_x = -\frac{(\omega_z + \omega)}{\omega_{xy}} f_z$ and $f_y = -(\omega_{xy} \tau)^{-1} f_z$. Note that, although we will not discuss these components further in this letter, they are an integral part of the model and crucial for the determination of f_z . An expression similar to Eqn. 5 has been used to describe Gilbert damping in a ferromagnet [8], and Eqn. 5 can be viewed as a generalization of that result. In resonance when $\omega_z = -\omega$, the universal value $\hbar \omega$ is achieved when $\omega_{xy} \tau \gg 1$. Hereafter, we will focus on the more general case, where a finite $(\omega + \omega_z)$ strongly damps the spin accumulation [14]. Then, in the limit of long spin relaxation times such that $\tau^{-1} \ll (\omega_{xy}, \omega_z + \omega)$, the prefactor reduces to $(1 + (\frac{\omega_z + \omega}{\omega_{xy}})^2)^{-1}$ and is independent of τ ; f_z will then attain the universal value of $\hbar \omega$ only when $\omega_{xy} \gg (\omega_z + \omega)$.

We now turn to discussing the hybrid system of Fig. 1b. The rotating field is only applied in the space $x < 0$ (the "pumped" region I), whereas for $x > 0$ there are no magnetic fields (region II). We will consider two physical realizations for the pumped region I: spin pumping in a non-magnetic material with an applied, rotating magnetic field of magnitude $B_{xy} = 10$ mT, and $B_z = 0$ (both region I and II are non-magnetic conductors); and a model of a weak ferromagnet with $B_{xy} = 1$ T and $B_z = 100$ T representing the exchange interaction corresponding to small angle magnetization precession around an easy axis (region II is a non-magnetic conductor). This exchange interaction approach, in which the magnetization of the localized d-moments couples to the s-electrons and polarizes them, has been used in early spin-pumping theory [15] (which was applied to transmission electron spin resonance), and more recently in the mean-field-theory description of Gilbert damping [8].

The system is solved as a boundary-value problem governed by Eqn. 4 with the diffusion term reinstated:

$$\lambda^2 \nabla^2 \vec{f} = \vec{f} + (\vec{\omega}_B + \vec{\omega}) \tau \times \vec{f} + \hbar \vec{\omega} \times \vec{\omega}_B \tau. \quad (6)$$

This equation can either be solved numerically directly, for instance using a finite element method solver, or by using the boundary conditions to determine the six constants of integration (for each region) of the general solution [16]. For simplicity in our model system we have




FIG. 2: The spin accumulation $f_z/\hbar\omega$ as a function of x/λ away from the interface, for τ ranging in decades from 100 ps to 10 s. Both regions are unbounded. For (a), $B_{xy} = 10$ mT and $B_z = 0$, for (b) $B_{xy} = 1$ T and $B_z = 100$ T.

assumed that the spin diffusion lengths, times and diffusion constants are the same for both regions. With these assumptions the boundary conditions at the interface reduce to continuity of $\vec{f}$ and $\nabla\vec{f}$ at $x = 0$. When the regions are unbounded, the outer boundary conditions are that in region I, the bulk result (Eqn. 5 and below) is recovered; in region II, $\vec{f}(x \gg 0) = 0$. When the regions are bounded, the boundary condition is that there is no leakage of spin current, so $\nabla\vec{f} = 0$ at the boundaries. In the calculations the diffusion constant has been fixed at $D = 0.002 \text{ m}^2/\text{s}$ (such as for Al) so the spin diffusion length λ scales as $\sqrt{\tau}$, and $\omega = 10$ GHz.

We first consider the case where both regions are unbounded. Fig. 2 shows the results of the calculation for the two different field pumping regimes, plotting the spatial dependence of $f_z/\hbar\omega$ near the interface for values of τ ranging over many decades. The most striking feature in Fig. 2 is that the spin accumulation at the interface is strongly enhanced over the bulk, uniform system value (at $x \ll 0$), and in the limit of very large τ achieves the universal value $\hbar\omega$, similar to the results of Ref. [2]. The calculations seem to describe a spin current injected at the interface which diffuses into both regions, even though the pumping occurs uniformly throughout region I. The enhancement of the spin accumulation at the interface is the primary result of this letter. In the unbounded system this effect is universal only for unphysically large τ , so we now try bounding the system




FIG. 3: Spin accumulation at the interface for the symmetrically bounded system plotted as a function of the normalized bounding length L/λ_ω , where $\lambda_\omega = \sqrt{2\pi D/\omega} \simeq 1.1 \mu\text{m}$, for τ ranging in decades from 10^{-10} (bottom curve) to 10 s (top curve).

in order to optimize the effect. In what follows we will focus only on the weak ferromagnet model, however most of the results apply generally to both systems. Fig. 3 shows the spin accumulation at the interface obtained for symmetric bounding of the regions at $x = \pm L$ as a function of L/λ_ω , over decades of τ from 10^{-10} to 10 s. The spin accumulation is optimized when the sample dimension $L \simeq 0.6\lambda_\omega$, independent of τ . Fig. 4 shows the dependence of the spin accumulation on the bounding length, calculated under various bounding conditions for a particular $\tau = 10^{-7}$ s. Bounding only region II yields optimal spin accumulation for L/λ_ω close to one, however upon bounding only the pumped region the spin accumulation peaks at a smaller length scale, corresponding to $L \simeq \lambda_B$. Bounding region II at $L_2 = 0.6\lambda_\omega$ and allowing L_1 to vary yields a peak at $L_1 \simeq 0.6\lambda_B$. The spin accumulation obtained for different specific bounding lengths as a function of τ is summarized in Fig. 5. The main message from this figure is that by engineering the sample dimensions it is possible to achieve universality at much lower τ .

A qualitative explanation for the interface enhancement is the averaging effect region II has on the x- and y-components of the spin accumulation. These components diffuse into region II but can diffuse back after some delay time. The back-flow will have a distribution of spin angles with respect to the rotating field in region I that tends to average out their influence, an effect similar to increasing the spin relaxation rate τ_{xy}^{-1} for the x- and y-components of the spin accumulation. However f_z does not experience such an averaging effect. In a simple model of spin accumulation in a uniform system it can be shown that reducing τ_{xy} while staying in the limit $\omega_{xy}^2 \tau_{xy} \tau_z \gg 1$ leads to an enhancement of the spin accumulation [16].




FIG. 4: Spin accumulation at the interface vs the normalized bounding length L/λ_ω , calculated with $\tau = 10^{-7}$ s under various bounding conditions indicated as L_1, L_2 in the figure. The largest enhancement is obtained for the $L, 0.6\lambda_\omega$ case with $L \simeq 0.6\lambda_B = 0.6\sqrt{2\pi D/\omega_B} = 15$ nm.




FIG. 5: The spin accumulation at the interface vs τ for specific bounding lengths indicated as L_1, L_2 (curves A through E). The open circles are based on Ref. [2] with $\eta = 10^{-5} \frac{1}{\sqrt{\tau}} \tanh \sqrt{\frac{2\pi}{\omega\tau}}$ (see text for details).

In the interface scattering matrix model [2], the spin accumulation close to the interface is given as $f_z = \hbar\omega \frac{\sin^2 \theta}{\sin^2 \theta + \eta}$, where θ is the precession cone angle and $\eta = (\tau_i/\tau) \tanh(L/\lambda)/(L/\lambda)$ is a reduction factor which depends on the normal metal properties and the spin-injection rate τ_i which is inversely proportional to the mixing conductance $g_{\uparrow\downarrow}$. In the limit of very large τ and λ , the reduction factor goes to zero and the cone angle dependence factors out, leaving the maximum spin accumulation $f_z = \hbar\omega$. The authors restrict their analysis to the case where the ferromagnet thickness should be less than the spin diffusion length. To compare to their result we take $\theta \simeq \frac{B_{xy}}{B_z} = \frac{1}{100}$ and evaluate η for $L = \lambda_\omega$, and find that η depends on τ as $\eta \propto \frac{1}{\sqrt{\tau}} \tanh \sqrt{\frac{2\pi}{\omega\tau}}$. This functional dependence on τ agrees well with curves D and

E, as shown for the fit to curve E (with 10^{-5} as the fitting parameter for η), even though our model is derived for an entirely different regime.

Our model of spin pumping in a non-magnetic region by a rotating magnetic field can be applied quite generally to diffusive systems, such as metals and semiconductors. Considering a semiconductor with τ as large as 100 ns and experimentally quite accessible $\omega = 10^{12}$ Hz, then a spin accumulation close to the universal result of $\hbar\omega$ corresponding to 700 μ V can be achieved even in the non-resonant case. The measurement of small spin accumulation signals due to the spin Hall effect in semiconductors by Kerr rotation microscopy has recently been demonstrated [17]. The spin pumping-induced spin accumulation and its interface enhancement that we have described should be measurable in similar systems.

We acknowledge useful discussions with G. E. W. Bauer and M. Costache. Support has been provided by the Stichting Fundamenteel Onderzoek der Materie (FOM) and the RTN Spintronics Network.

* Now at Unité Mixte de Physique CNRS/Thales, Route départementale 128, 91767 Palaiseau Cedex, France.

- [1] L. Berger, Phys. Rev. B **59**, 11465 (1999).
- [2] A. Brataas, Y. Tserkovnyak, G. E. W. Bauer, and B. I. Halperin, Phys. Rev. B **66**, 060404(R) (2002).
- [3] Spin pumping in systems with (quantum) confinement is discussed in P. Sharma, Science **307**, 531 (2005).
- [4] A. Brataas, Y. V. Nazarov, and G. E. W. Bauer, Phys. Rev. Lett. **84**, 2481 (2000).
- [5] B. Heinrich, Y. Tserkovnyak, G. Woltersdorf, A. Brataas, R. Urban, and G. E. W. Bauer, Phys. Rev. Lett. **90**, 187601 (2003); R. Urban, G. Woltersdorf, and B. Heinrich, Phys. Rev. Lett. **87**, 217204 (2001).
- [6] S. Mizukami, Y. Ando, and T. Miyazaki, J. Magn. Magn. Mater. **239**, 42 (2002).
- [7] Y. Tserkovnyak, A. Brataas, and G. E. W. Bauer, Phys. Rev. Lett. **88**, 117601 (2002).
- [8] Y. Tserkovnyak, G. A. Fiete, and B. I. Halperin, Appl. Phys. Lett. **84**, 5234 (2004).
- [9] A spin accumulation $\tilde{f}(x, t)$ is in general not directly proportional to the magnetization. However, f_z in Eqn. 5, as a steady-state quantity, is associated with a contribution to the magnetization of $\frac{1}{2}\mu_B N(E_F) f_z$, where $N(E_F)$ is the density of states at the Fermi level.
- [10] D. H. Hernando, Y. V. Nazarov, A. Brataas, and G. E. W. Bauer, Phys. Rev. B **62**, 5700 (2000).
- [11] M. Johnson and R. H. Silsbee, Phys. Rev. B **37**, 5312 (1988).
- [12] F. J. Jedema, H. B. Heersche, A. T. Filip, J. J. A. Baselmans, and B. J. van Wees, Nature **416**, 713 (2002).
- [13] Quantum-mechanically this cannot be fully justified, however we work in a strictly classical ensemble limit such that temporal and spatial coherence and interference effects can be neglected.
- [14] It is difficult to realize the resonant case experimentally when pumping from a ferromagnet. The resonant case

is treated elsewhere for pumping from a non-magnetic material [16].

- [15] R. H. Silsbee, A. Janossy, and P. Monod, Phys. Rev. B **19**, 4382 (1979).
- [16] S. M. Watts *et al.*, in preparation.
- [17] Y. K. Kato, R. C. Myers, A. C. Gossard, and D. D. Awschalom, Science **306**, 5703 (2004).